




2103 W. Houston St.
Sherman, Texas 75092
Phone : (903)-893-5561
Fax : (903)-893-6598
Email : lange.vet.hospital@hotmail.com



Dr. Grayson Riddels
Dr. Samantha Riddels
Dr. William T. Lange
Dr. Leah Eaton

Puppy Visit Schedule

1st Visit : 6-9 Weeks _____

- 1st DHPP (Distemper, Adenovirus, Parainfluenza, Parvovirus)
- Fecal Check
- Start Parasite Prevention
- Pet Insurance Registration

2nd Visit : 9-12 Weeks _____

- 2nd DHPP (Distemper, Adenovirus, Parainfluenza, Parvovirus)
- Recheck Fecal
- Refill Prevention

3rd Visit : 12-15 Weeks _____

- DHLPP (Distemper, Lepto, Adenovirus, Parainfluenza, Parvovirus)
- Oral Bordetella
- Refill Prevention

4th Visit : 16-18 Weeks _____

- DHLPP (Distemper, Lepto, Adenovirus, Parainfluenza, Parvovirus)
- 1YR Rabies
- Refill Prevention
- Schedule Spay/Neuter Surgery (Microchip can be done at time of surgery)

Puppy boosters should be given every 3-4 weeks up to 16 weeks regardless of how many previous vaccines they have received

DHPP - This is a 4-way vaccine which is recommended for every dog. Distemper is a severe and sometimes deadly virus that can affect the entire body, including the brain. Parvo is a dangerous virus which causes severe intestinal disease and dehydration. Puppies are given a series of DHPP vaccinations, 3-4 weeks apart, usually starting at 6 weeks old and ending at 14-16 weeks of age. The DHPP vaccine is boosted 1 year later and is then recommended annually thereafter.

Bordetella - Bordetella causes upper respiratory disease which can sometimes lead to pneumonia. This generally presents itself as a persistent cough or tracheal bronchitis. The vaccine is recommended for dogs that come into contact with other dogs (grooming, boarding, park, etc...). It is given to puppies at 12-15 weeks of age and then yearly thereafter.

Lepto - Dogs can get this bacterial disease through contact with the urine of a variety of animals, including rodents. It can lead to kidney and liver failure and can be passed on to people. The vaccine is recommended for dogs who may drink from or wade in standing pools of water or are prone to chasing small rodents.

Rabies - Rabies is an invariably lethal disease which causes abnormal and sometimes aggressive behavior. It can be passed on to people. Rabies vaccination of dogs is required by Texas law at 16 weeks of age, 1 year later, and then every 3 years thereafter.

Fecal - We recommend performing a fecal test at their first visit to check for intestinal parasites and then rechecking a fecal as needed. Fecal tests are recommended annually thereafter as well as year round intestinal parasite prevention.

Heartworm Test - We recommend a heartworm test annually for all pets as well as monthly heartworm prevention.